

Reproductive Technology Evaluation and Treatment of Infertility

Guidelines for Catholic Couples¹

Technologies Compatible with Catholic Teachings

1. Observation of the naturally occurring sign(s) of fertility (Natural Family Planning). Time intercourse on the days of presumed (potential) fertility for at least six months before proceeding to medical interventions.²
2. General medical evaluation of both spouses for infertility.
3. Post-coital test to assess sperm number and viability in “fertile type” mucus. These tests are undertaken after normal intercourse.
4. Appropriate evaluation and treatment of male factor deficiency. Seminal fluid samples can be obtained from a non-lubricated, perforated condom after normal intercourse.
5. Assessment of uterine and tubal structural competence by imaging techniques (e.g., ultrasound, hysterosalpingogram, etc.).
6. Appropriate medical treatment of ovulatory dysfunction.
7. Appropriate (usually surgical) correction of mechanical blocks to tubal patency (the state of being open).

Reproductive Technology Under Discussion

Some reproductive technologies continue to be under study. Although the Church has not officially condemned them, Catholics are urged to learn about this discussion and in conscience determine if they should refuse or use them.

Intrauterine insemination (IUI) of “licitly obtained” (normal intercourse) but technologically prepared semen sample (washed, and reinserted, etc.).

- *Comment:* The probable objection of the Church is to the “third-party” involvement that happens between the marital act and conception. In other words—*Is this an “intrusive intervention” between conjugal relations and conception?* The main ethical point regarding the Church’s condemnation of artificial insemination is that a third-party intrudes in the conjugal act.

Reproductive Technologies in Disagreement with Catholic Teachings (*wrong*)

1. Obtaining a sample of seminal fluid by masturbation.
2. Artificial insemination by a non-spouse (AID)
3. Artificial insemination by the husband (AIH) if the sample is obtained and handled by non-licit means (masturbated specimen).

4. Any use of gametes (egg and sperm) from outside the marital relationship.
5. Any use of a gestational carrier (“surrogate mother”).
6. In vitro fertilization (IVF), zygote intra-fallopian transfer (ZIFT), and intracytoplasmic sperm injection (ICSI), ovum donation, “surrogate” uterus.
7. Freezing embryos.

“How do I know when a reproductive technology is morally right?”

The rule of thumb is

Procedures that remove obstacles to natural fertility and which assist marital intercourse in reaching its procreative potential are morally acceptable.

Procedures which add a “third party” into the act of conception or gestation, or which substitute a laboratory procedure for intercourse, are not morally acceptable.

NOTES

1. These guidelines are drawn from the document [Donum vitae](#) (1987). See also [Dignitas personae](#) (2008).
 2. Some women may have to wait longer for their fertility signs to appear due to the effects of use of chemical contraceptives (e.g., the pill, shots, patches, etc.).
-

Definitions

AID (Artificial insemination with donor's sperm)

Donor sperm is inserted into a cup and placed over the woman's cervix.

AIH (Artificial insemination with husband's sperm)

A husband's sperm is inserted into a cup which is placed over his wife's cervix.

ICSI (Intra-Cytoplasmic Sperm Injection)

When men have low sperm counts or other problems, such as blocked ducts, spermatozoa can be obtained either by masturbation or, in the absence of a normal ductal system, by needle aspiration from the epididymis or even from the testis itself. A single sperm is then injected through the membrane of the ovum and the embryo cultured in the laboratory until it reaches the 8-16 cell stage, when it is inserted into the uterine cavity.

Because the "natural selection" which occurs when sperm enter through the cervical mucus is excluded by this procedure, a number of birth defects have been recorded when conception was effected by ICSI.

IVF (In Vitro Fertilization)

Conception occurs outside the body—"in a glass."

Ordinarily, the woman is treated with hormones to stop her natural cycle and stimulated to ripen a number of ova. The ova are harvested from the follicle with a needle under ultrasonic guidance. The needle is inserted either through the vagina or abdomen. Ova are incubated in the laboratory with a carefully washed and adjusted specimen of semen to allow fertilization. Prior to implantation in the woman's uterus, embryos are examined in order to select the "best." Sometimes, one cell is removed for genetic testing. To date, visual inspection of the embryos has been totally unrelated to their subsequent course—health or otherwise. Usually at least two embryos are implanted; in some centers, as many as four are implanted with the hope of getting at least one live baby. At times, three or four embryos thrive. Some clinics then offer the mother "embryo reduction" (or "fetal reduction"; this is selective abortion) to allow only one or two preborn babies (fetuses) to develop further.

Because the endometrium is considerably changed by the stimulation of ovaries to produce eggs, it is the practice in some centers to freeze the embryos and to implant them in a subsequent natural cycle. Success rates vary considerably due to the age of the woman, the health of the uterus, the quality of semen, etc. The disposition of frozen embryos varies with the wishes of the parents. "Spare embryos" may either be preserved, donated to other women or to researchers, or destroyed.

IUI (intrauterine insemination) of "licitly obtained" (normal intercourse) but technologically prepared semen sample.

The sperm are collected from a perforated condom after normal intercourse, washed, and then injected into the uterine cavity, bypassing the cervix to avoid "hostile" mucus. Cervical mucus hostility is an immunological reaction brought about by several known, and some unknown factors. A postcoital test would find no living sperm in mucus during the fertile phase. Other treatments for cervical mucus hostility include abstinence for two years to allow the antibodies to diminish or disappear, or the use of condoms (not acceptable for Catholics). Various treatments with steroids have been tried without much success.

ZIFT (Zygote Intra-Fallopian Transfer)

Ova and sperm are obtained analogously to IVF, but the zygote, that is the newly fertilized embryo, is immediately transferred into the woman's tube with a catheter threaded through the uterus. This does not allow examination of the embryos as it would for IVF. The live birth rate is similar to IVF.



Gratitude is extended to the original author of this text, medical gynecologist, NFP pioneer, and religious sister, Hanna Klaus, MD (Sr. Miriam Paul). This resource has been updated in consultation with staff at the National Catholic Bioethics Center (NCBC).

Copyright © 2025, United States Conference of Catholic Bishops. All rights reserved. Permission is granted to reproduce in whole or in part, in print and/or electronically, with the following statement: Last Name, First Name of Author, "Title," NFPP/US Conference of Catholic Bishops, Washington, DC: USCCB, 2025. Used with permission.
